

LONG TERM CARE SURVEY

SURVEY AREA	OBSERVATION	INTERVIEWING	RECORD REVIEW	EVALUATION FACTORS	CROSS REFERENCE															
F196 (cont'd)			<p>Example: 120 lbs/2.2 lbs. = 54.5 kg (55 kgs) 55 kg x 30 cc = 1,650 cc/day</p> <p>Note: Isotonic Standard Tube Feeding = Approximately 80% water.</p> <p><u>Amputation % of Body Weight</u></p> <p>Leg 20% Below Knees 10% Arm 6% At Elbow 3.6%</p> <p><u>Suggested Standards for Evaluating Significance of Weight Loss</u></p> <p>% of body weight loss</p> <table border="0"> <tr> <td>Inter- val</td> <td>Significant Loss</td> <td>Severe Loss</td> </tr> <tr> <td>1 week</td> <td>1-2%</td> <td>2%</td> </tr> <tr> <td>1 month</td> <td>5%</td> <td>5%</td> </tr> <tr> <td>3 months</td> <td>7 1/2%</td> <td>7 1/2%</td> </tr> <tr> <td>6 months</td> <td>10%</td> <td>10%</td> </tr> </table> <p>From Blackburn, et al: "Nutritional and Metabolic Assessment of the Hospitalized Patient: JPEN vol. 1, 1977.</p>	Inter- val	Significant Loss	Severe Loss	1 week	1-2%	2%	1 month	5%	5%	3 months	7 1/2%	7 1/2%	6 months	10%	10%		
Inter- val	Significant Loss	Severe Loss																		
1 week	1-2%	2%																		
1 month	5%	5%																		
3 months	7 1/2%	7 1/2%																		
6 months	10%	10%																		